

Constructing Calabi-Yau Metrics From Hyperkähler Spaces

H. Lü[†], Yi Pang[‡] and Zhao-Long Wang^{*}

[†]*China Economics and Management Academy*

Central University of Finance and Economics, Beijing, 100081, China

[‡]*Key Laboratory of Frontiers in Theoretical Physics*

Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100190

^{*}*Interdisciplinary Center for Theoretical Study,*

University of Science and Technology of China, Hefei, Anhui 230026, China

ABSTRACT

Recently, a metric construction for the Calabi-Yau 3-folds from a four-dimensional hyperkähler space by adding a complex line bundle was proposed. We extend the construction by adding a $U(1)$ factor to the holomorphic $(3, 0)$ -form, and obtain the explicit formalism for a generic hyperkähler base. We find that a discrete choice arises: the $U(1)$ factor can either depend solely on the fibre coordinates or vanish. In each case, the metric is determined by one differential equation for the modified Kähler potential. As explicit examples, we obtain the generalized resolutions (up to orbifold singularity) of the cone of the Einstein-Sasaki spaces $Y^{p,q}$. We also obtain a large class of new singular CY3 metrics with $SU(2) \times U(1)$ or $SU(2) \times U(1)^2$ isometries.

1 Introduction

Six-dimensional Calabi-Yau manifolds (CY3) [1, 2, 3] play an important role in string theory, since they provide natural internal compactifying spaces, giving rise to four-dimensional theories that preserve one quarter of the ten-dimensional supersymmetry. String compactification on the CY3 has been studied mainly based on their topological properties. This is because although it was demonstrated that Ricci-flat and complete metrics exist on the compact CY3 manifolds [3], one does not expect to see an explicit one, aside from the flat metric. This can be seen from the Killing vector analysis. A Killing vector K_i satisfies the equation

$$-\square K_i - R_{ij}K^j = 0. \quad (1)$$

Multiply by K^i and integrate over the manifold. For a compact manifold, integration by parts on the first term gives no boundary contribution, and hence one concludes

$$\int_{\mathcal{M}} (|\nabla_i K_j|^2 - R_{ij}K^i K^j) = 0. \quad (2)$$

For the Ricci-flat CY3 metrics, it must be that $\nabla_i K_j = 0$ pointwise everywhere in the manifold. Leaving aside the trivial possibility that there are flat S^1 factors, such a covariantly-constant vector will not exist. Therefore there can be no Killing vectors in a non-flat Ricci-flat compact manifold. Without Killing vectors, there are no continuous symmetries. And without the simplifications that result from supposing that a metric has continuous symmetries, it is essentially hopeless to solve the Einstein equations.

Thus explicit metrics on the CY3 with Killing vectors are necessarily non-compact and/or singular. In fact, in string theory compactification, the CY3 spaces can develop singularities at limiting values of their modulus parameters, where additional massless four-dimensional states will emerge. The simplest example of such a singular metric is the conifold, which is the Ricci-flat metric on the cone over the homogeneous Einstein-Sasaki space $T^{1,1} = (S^3 \times S^3)/U(1)_{\text{diag}}$. The metric is singular at the vertex of the cone, and as the moduli are moved slightly away from the singular limit, the metric near to the previous conifold point is then smoothed out. It was shown [4] that there are two possible ways of smoothing out the conifold: in one of which the vertex is blown up to an S^2 ; in the other, it is blown up to an S^3 .

Not so long ago, it was demonstrated by explicit construction that there exist an infinite number of Einstein-Sasaki metrics with the toric $U(1)^3$ isometry, called $Y^{p,q}$ [5] and $L^{p,q,r}$ [6, 7]. These provide an infinite number of generalized (toric) conifolds, some of which

can be smoothed out completely [8, 9, 10] or up to an orbifold singularity [9, 10]. The construction can be generalized to arbitrary $2n$ dimensions, since it turns out that the local metrics can be obtained directly from the BPS limits [11, 6, 12] of the Euclidean version of the general Kerr-AdS black hole [13, 14, 15] and Kerr-AdS-NUT solutions [12] in arbitrary dimensions: the odd dimensional ones give rise to Einstein-Sasaki spaces whilst the even dimensional ones give rise to Calabi-Yau metrics.

The same Killing vector analysis applies to the higher dimensional manifolds with reduced holonomy and one does not expect to see an explicit metric on compact spaces. Examples of complete metrics on non-compact manifolds in higher dimensions with G_2 and $spin(7)$ holonomy were constructed in the later 1980's [16, 17, 18, 19]. Inspired by the AdS/CFT correspondence and applications in M-theory compactification, large classes of explicit metrics on G_2 and $spin(7)$ holonomy spaces have been constructed and their applications in string and M-theory have been discussed [20]-[39].

Although a large number of Calabi-Yau metrics have been constructed, an organizing principle is still lacking, since many of these metrics are discovered serendipitously, or constructed indirectly through the BPS limit of the known Kerr-AdS-NUT solutions. Recently, a new technique was developed in [40] for constructing CY3 metrics, generalizing the construction of the D6-brane wrapping on a two-cycle of a four-dimensional hyperkähler space [41]. The essence of the construction is to build a CY3 metric from a hyperkähler one by adding a complex line bundle. This follows the same line of constructing $(2n+2)$ -dimensional Einstein-Kähler spaces from $(2n)$ -dimensional ones [17]. Differently, in the new construction, the Kähler potential in four dimensions is allowed to be modified by an arbitrary function G . However, the proof of the existence of such CY3 metrics were presented for the $\mathbb{R}^4$ base only. What is curious is that the metrics with the asymptotic structure of cones over Einstein-Sasaki spaces, such as the $\mathbb{R}^6$ or conifolds, are absent from the construction presented in [40].

In section 2, we extend this construction by considering a generic hyperkähler base. Furthermore, we find that the ansatz for the holomorphic $(3,0)$ -form presented in [40] can be supplemented with a $U(1)$ factor. This allows us to construct a much wider class of solutions including ones that have asymptotic cones over Einstein-Sasaki spaces. There are two discrete possibilities for the $U(1)$ factor. One is that it is dependent solely on the fibre coordinates. In this case the equations for the CY3 are reduced to one differential equation for the modified Kähler potential G . The other possibility is that the $U(1)$ factor vanishes, for which the metric is determined by a differential equation which is the singular limiting

case of the previous one. In section 3, we consider the simplest $\mathbb{R}^4$ base and obtain the CY3 metrics that describe resolutions of the cone over the $Y^{p,q}$ spaces, when the $U(1)$ factor in the $(3,0)$ -form is non-vanishing. For the case with vanishing $U(1)$ factor, we obtained a large class of new singular metrics. In section 4, we consider triaxial basis for the hyperkähler spaces that preserve the $SU(2)$ isometry. Again solutions with or without the $U(1)$ factors were obtained. We conclude our paper in section 5. In appendices A and B, we present some complicate formulae and detailed derivations.

2 The construction

2.1 The ansatz

In this section, we review and then extend significantly the metric construction for the CY3 from $D = 4$ hyperkähler spaces, proposed in [40]. Let us consider a generic hyperkähler space in four dimensions with the complex coordinates $z^i, \bar{z}^i$ ($i = 1, 2$) and the Kähler potential $K_0(z^i, \bar{z}^i)$. The metric is given by

$$ds^2 = 2\tilde{g}_{i\bar{j}}dz^i d\bar{z}^j, \quad \tilde{g}_{i\bar{j}} = \frac{1}{2}\partial_i\partial_{\bar{j}}K_0 = \tilde{g}_{\bar{j}i}. \quad (3)$$

Since it is Ricci flat, the Ricci form $\mathcal{R}^{(1,1)}$ vanishes, *i.e.*

$$\mathcal{R}^{(1,1)} = i\tilde{\partial}\bar{\partial}\log\sqrt{V} = 0. \quad (4)$$

Here $V \equiv \det(\tilde{g}_{i\bar{j}})^2$ is the volume factor and $\tilde{\partial}$ and $\bar{\partial}$ are the Dolbeault 1-form differential operators defined by

$$\tilde{\partial} \equiv dz^i \partial_{z^i}, \quad \bar{\partial} \equiv d\bar{z}^i \partial_{\bar{z}^i}. \quad (5)$$

The equation (4) implies that $\log V$ is the real part of a holomorphic function, or equivalently, V can be the norm of a holomorphic function. The choice for the complex coordinates is not unique since we can always make a holomorphic coordinate transformation $z^i \rightarrow z'^i = f^i(z^j)$. Under such a transformation, the volume factor transforms as

$$V \rightarrow |T|^{-4}V, \quad (6)$$

where

$$T = \det \left[\frac{\partial(f^1, f^2)}{\partial(z^1, z^2)} \right]. \quad (7)$$

It is easy to see that T can be any holomorphic function. Thus we can always set $V = 1$ by choosing appropriate complex coordinates. We shall do this for later convenience.

We now consider a complex vielbein basis of the hyperkähler space, which is given by $\tilde{\epsilon}^a, \bar{\tilde{\epsilon}}^a$ ($a = 1, 2$), then the corresponding metric, Kähler form and holomorphic $(2, 0)$ -form are given by

$$\begin{aligned} ds^2 &= \tilde{\epsilon}^1 \bar{\tilde{\epsilon}}^1 + \tilde{\epsilon}^2 \bar{\tilde{\epsilon}}^2, \\ \tilde{J}^{(1,1)} &= \frac{i}{2}(\tilde{\epsilon}^1 \wedge \bar{\tilde{\epsilon}}^1 + \tilde{\epsilon}^2 \wedge \bar{\tilde{\epsilon}}^2), \\ \tilde{\Omega}^{(2,0)} &= \tilde{\epsilon}^1 \wedge \tilde{\epsilon}^2. \end{aligned} \quad (8)$$

According to the properties of hyperkähler spaces, both $\tilde{J}$ and $\tilde{\Omega}$ are closed, *i.e.* $d_4 \tilde{J} = d_4 \tilde{\Omega} = 0$. One can now use this structure to construct a CY3. The metric ansatz is given by [40]

$$\begin{aligned} ds^2 &= ds_4^2 + h^2 dy^2 + h^{-2}(d\alpha + A)^2 \\ &= (\delta_{a\bar{b}} + G_{a\bar{b}})\tilde{\epsilon}^a \bar{\tilde{\epsilon}}^b + h^2 dy^2 + h^{-2}(d\alpha + A)^2. \end{aligned} \quad (9)$$

The metric components are the functions of the $y, z^i, \bar{z}^i$ coordinates, but are independent of the coordinate α , which is a manifest Killing direction. Thus the metric ansatz assumes at least one Killing direction. The functions appearing in the ds_4^2 part are defined by

$$\tilde{\partial} \bar{\partial} G = dz^i \wedge d\bar{z}^j \partial_i \partial_{\bar{j}} G = \tilde{\epsilon}^a \wedge \bar{\tilde{\epsilon}}^b \tilde{\epsilon}_a^i \bar{\tilde{\epsilon}}_b^j \partial_i \partial_{\bar{j}} G = G_{a\bar{b}} \tilde{\epsilon}^a \wedge \bar{\tilde{\epsilon}}^b, \quad (10)$$

where $\tilde{\epsilon}_a$ is the inverse complex vielbein. Note that if we replace G by K_0 in (10), we have $\delta_{a\bar{b}}$ instead of $G_{a\bar{b}}$. Thus the ds_4^2 in (9) is obtained by deforming the original Kähler potential $K_0(z^i, \bar{z}^i)$ to $K_0(z^i, \bar{z}^i) + G(y, z^i, \bar{z}^i)$.

The ds_4^2 can be diagonalized by a local $SU(2)$ transformation $U_a{}^b(z^i, \bar{z}^i)$, namely

$$U \begin{pmatrix} 1 + G_{1\bar{1}} & G_{1\bar{2}} \\ G_{2\bar{1}} & 1 + G_{2\bar{2}} \end{pmatrix} U^\dagger = \begin{pmatrix} \lambda_1(z^i, \bar{z}^i) & 0 \\ 0 & \lambda_2(z^i, \bar{z}^i) \end{pmatrix}. \quad (11)$$

We further suppose that the complex structure of the hyperkähler base is part of complex structure of the CY3. This implies that the complex vielbein of the CY3 is given by

$$\epsilon^1 = e^{i\frac{\kappa}{2}} \sqrt{\lambda_1} \tilde{\epsilon}^a (U^\dagger)_a{}^1, \quad \epsilon^2 = e^{i\frac{\kappa}{2}} \sqrt{\lambda_2} \tilde{\epsilon}^a (U^\dagger)_a{}^2, \quad \epsilon^3 = h dy + i h^{-1} (d\alpha + A). \quad (12)$$

where $\kappa = \kappa(\alpha, y, z^i, \bar{z}^i)$ is a real function. Correspondingly, the Kähler form and the $(3, 0)$ -form for the CY3 are given by

$$J^{(1,1)} = \frac{i}{2}(\epsilon^1 \wedge \bar{\epsilon}^1 + \epsilon^2 \wedge \bar{\epsilon}^2 + \epsilon^3 \wedge \bar{\epsilon}^3) = \frac{i}{2}(\delta_{a\bar{b}} + G_{a\bar{b}})\tilde{\epsilon}^a \wedge \bar{\tilde{\epsilon}}^b + dy \wedge (d\alpha + A), \quad (13)$$

$$\Omega^{(3,0)} = \epsilon^1 \wedge \epsilon^2 \wedge \epsilon^3 = f e^{i\kappa} \tilde{\epsilon}^1 \wedge \tilde{\epsilon}^2 \wedge (h dy + i h^{-1} (d\alpha + A)), \quad (14)$$

where

$$f = \sqrt{\lambda_1 \lambda_2} = \sqrt{\det(\delta_{a\bar{b}} + G_{a\bar{b}})} = \sqrt{1 + G_{1\bar{1}} + G_{2\bar{2}} + G_{1\bar{1}}G_{2\bar{2}} - G_{1\bar{2}}G_{2\bar{1}}}. \quad (15)$$

Note that comparing with the ansatz in [40], we have introduced a $U(1)$ factor $e^{i\kappa}$ for the holomorphic $(3,0)$ -form. This factor turns out to be crucial for constructing metrics with asymptotic cones over Einstein-Sasaki spaces. Although the metric is independent of the coordinate α , this $U(1)$ factor can be.

The requirement that the metric (9) be Calabi-Yau becomes the requirement that the above Kähler form and $(3,0)$ -form are both closed. Note that

$$\begin{aligned} dJ &= \frac{i}{2} \partial_y G_{a\bar{b}} dy \wedge \tilde{\epsilon}^a \wedge \bar{\epsilon}^b - dy \wedge d_4 A = \frac{i}{2} dy \wedge \tilde{\partial} \bar{\partial} \partial_y G - dy \wedge d_4 A \\ &= -\frac{i}{4} dy \wedge (\tilde{\partial} + \bar{\partial})(\partial - \bar{\partial}) \partial_y G - dy \wedge d_4 A = -\frac{i}{4} dy \wedge d_4 (\partial - \bar{\partial}) \partial_y G - dy \wedge d_4 A, \end{aligned} \quad (16)$$

then $dJ = 0$ implies that

$$A = -\frac{i}{4} (\tilde{\partial} - \bar{\partial}) \partial_y G + \lambda(y, z_i, \bar{z}_i) dy, \quad (17)$$

up to some pure gauge terms. Note that d_4 denotes an exterior derivative with respect to z^i and $\bar{z}^i$ only. The exterior derivative for the $(3,0)$ -form is given by

$$\begin{aligned} d\Omega &= \bar{\partial}(f e^{i\kappa} h) \wedge \tilde{\epsilon}^1 \wedge \tilde{\epsilon}^2 \wedge dy + i f e^{i\kappa} h \partial_\alpha \kappa d\alpha \wedge \tilde{\epsilon}^1 \wedge \tilde{\epsilon}^2 \wedge dy \\ &\quad + i \bar{\partial}(f e^{i\kappa} h^{-1}) \wedge \tilde{\epsilon}^1 \wedge \tilde{\epsilon}^2 \wedge \left(d\alpha + \frac{i}{4} \bar{\partial}(\partial_y G) + \lambda dy \right) \\ &\quad + i \partial_y (f e^{i\kappa} h^{-1}) dy \wedge \tilde{\epsilon}^1 \wedge \tilde{\epsilon}^2 \wedge \left(d\alpha + \frac{i}{4} \bar{\partial}(\partial_y G) \right) \\ &\quad - i f e^{i\kappa} h^{-1} \tilde{\epsilon}^1 \wedge \tilde{\epsilon}^2 \wedge dy \wedge \bar{\partial} \lambda - \frac{i}{4} f e^{i\kappa} h^{-1} dy \wedge \tilde{\epsilon}^1 \wedge \tilde{\epsilon}^2 \wedge \bar{\partial} \partial_y^2 G \\ &\quad - f e^{i\kappa} h^{-1} \partial_\alpha \kappa d\alpha \wedge \tilde{\epsilon}^1 \wedge \tilde{\epsilon}^2 \wedge \left(\frac{i}{4} \bar{\partial}(\partial_y G) + \lambda dy \right). \end{aligned} \quad (18)$$

The vanishing of the terms containing $d\alpha \wedge dy$ implies that

$$\partial_y (f h^{-1}) - f h \partial_\alpha \kappa = 0, \quad (19)$$

$$\partial_y \kappa - \lambda \partial_\alpha \kappa = 0. \quad (20)$$

Since by construction only κ can depend on α , it follows from (19) that we have

$$\kappa = \alpha \kappa_1(y, z^i, \bar{z}^i) + \kappa_0(y, z^i, \bar{z}^i). \quad (21)$$

Substituting it back to (20), we find

$$\kappa_1 \lambda = \partial_y \kappa_0, \quad \kappa = \alpha \kappa_1(z^i, \bar{z}^i) + \kappa_0(y, z^i, \bar{z}^i), \quad (22)$$

$$\partial_y g = \kappa_1 g^{-1} f^2, \quad (23)$$

where

$$g \equiv f h^{-1}. \quad (24)$$

The vanishing of the other terms containing $d\alpha$ implies that

$$4\bar{\epsilon}_a^\mu \partial_\mu (g e^{i\kappa}) + g e^{i\kappa} \kappa_1 \bar{\epsilon}_a^\mu \partial_\mu \partial_y G = 0, \quad (25)$$

from which, we find

$$\bar{\epsilon}_a^\mu \partial_\mu \kappa_1 = 0, \quad (26)$$

$$\bar{\epsilon}_a^\mu \partial_\mu (4 \log g + i 4 \kappa_0 + \kappa_1 \partial_y G) = 0. \quad (27)$$

Since κ_1 is real, (26) implies that κ_1 is a constant. It can be set to either 0, or 1, by rescaling the α coordinate. The vanishing of the rest terms implies that

$$\begin{aligned} & 4\bar{\epsilon}_a^\mu \partial_\mu (g^{-1} f^2 e^{i\kappa} + i g e^{i\kappa} \lambda) + \partial_y (g e^{i\kappa} \bar{\epsilon}_a^\mu \partial_\mu \partial_y G) = 0, \\ \Rightarrow & \bar{\epsilon}_a^\mu \partial_\mu (4 g^{-2} f^2 + i 4 \lambda + \partial_y^2 G) = 0, \end{aligned} \quad (28)$$

where we have used the equations (22), (23), (25) and (27).

2.2 Case I: $\kappa_1 = 1$

For $\kappa_1 = 1$, the equation (28) can be deduced from (23) and (27). Then the CY3 is determined by the following equations

$$g \partial_y g = 1 + G_{1\bar{1}} + G_{2\bar{2}} + G_{1\bar{1}} G_{2\bar{2}} - G_{1\bar{2}} G_{2\bar{1}}, \quad (29)$$

$$\bar{\epsilon}_a^\mu \partial_\mu (\partial_y G + 4 \log g + i 4 \kappa_0) = 0, \quad (30)$$

and the other quantities are then given by

$$\lambda = \partial_y \kappa_0, \quad \kappa = \alpha + \kappa_0(y, z^i, \bar{z}^i). \quad (31)$$

Note that a partial derivative of (30) with respect to y gives rise to (28). The equation (30) implies that we have $\partial_y G + 4 \log g + i 4 \kappa_0 = H(y, z^i)$, where

$$H \equiv U + i V \quad (32)$$

is any holomorphic function on the hyperkähler base. The real part $U = \partial_y G + 4 \log g$ satisfies

$$\bar{\epsilon}_a^\mu \bar{\epsilon}_b^\nu (\partial_\mu \partial_\nu - \Gamma_{\mu\nu}^\lambda \partial_\lambda) U = 0, \quad (33)$$

The imaginary part is then given by

$$4\kappa_0 = V = \int (\tilde{\partial} + \bar{\tilde{\partial}})V = -i \int (\tilde{\partial} - \bar{\tilde{\partial}})U. \quad (34)$$

However, note that $\partial_y G \rightarrow \partial_y G - U$ and $\alpha \rightarrow \alpha + \kappa_0$ are gauge transformations in our set up. Therefore, we can always set $H = 0$. Then we have $\kappa_0 = \lambda = 0$, and that

$$g = \exp\left(-\frac{1}{4}\partial_y G\right). \quad (35)$$

It follows from (29) that the system will be determined solely by the following basic equation for G

$$\partial_y \left[\exp\left(-\frac{1}{2}\partial_y G\right) \right] = 2(1 + G_{1\bar{1}} + G_{2\bar{2}} + G_{1\bar{1}}G_{2\bar{2}} - G_{1\bar{2}}G_{2\bar{1}}). \quad (36)$$

The $U(1)$ factor depends on the fibre coordinate α only.

2.3 Case II: $\kappa_1 = 0$

When $\kappa_1 = 0$, the CY3 is determined by the following equations

$$\bar{\epsilon}_a^\mu \partial_\mu (\log g + i\kappa_0) = 0, \quad (37)$$

$$\bar{\epsilon}_a^\mu \partial_\mu [\partial_y^2 G + 4g^{-2}(1 + G_{1\bar{1}} + G_{2\bar{2}} + G_{1\bar{1}}G_{2\bar{2}} - G_{1\bar{2}}G_{2\bar{1}}) + i4\lambda] = 0, \quad (38)$$

where $\kappa_0 = \kappa_0(z^i, \bar{z}^i)$ and $g = g(z^i, \bar{z}^i)$. It follows that both

$$g e^{i\kappa_0} = H_1(z^i) \quad (39)$$

and

$$\partial_y^2 G + 4g^{-2}(1 + G_{1\bar{1}} + G_{2\bar{2}} + G_{1\bar{1}}G_{2\bar{2}} - G_{1\bar{2}}G_{2\bar{1}}) + i4\lambda = H_2(y, z^i) = U + iV \quad (40)$$

are holomorphic functions on the hyperkähler base. Again, the gauge transformations $\partial_y^2 G \rightarrow \partial_y^2 G - U$ and $\alpha \rightarrow \alpha + \Lambda$ imply that we can always set $H_2 = 0$. Then we have $\lambda = 0$ and

$$\partial_y^2 G + 4g^{-2}(1 + G_{1\bar{1}} + G_{2\bar{2}} + G_{1\bar{1}}G_{2\bar{2}} - G_{1\bar{2}}G_{2\bar{1}}) = 0. \quad (41)$$

Furthermore, let us consider a holomorphic coordinate transformation $z^i \rightarrow \omega^i(z^j)$. It induces the following transformation on the four dimensional Kähler potential

$$G \rightarrow \tilde{G}(y, z^i, \bar{z}^i) = K_0(\omega^i, \bar{\omega}^i) - K_0(z^i, \bar{z}^i) + G(y, \omega^i, \bar{\omega}^i), \quad (42)$$

where K_0 is the Kähler potential for the hyperkähler base. If the holomorphic functions $\omega^i(z^j)$ satisfy

$$\det \left[\frac{\partial(z^1, z^2)}{\partial(\omega^1, \omega^2)} \right] = H_1(\omega^i), \quad (43)$$

it follows from (15) that the equation (41) becomes

$$\partial_y^2 \tilde{G} + 4(1 + \tilde{G}_{1\bar{1}} + \tilde{G}_{2\bar{2}} + \tilde{G}_{1\bar{1}}\tilde{G}_{2\bar{2}} - \tilde{G}_{1\bar{2}}\tilde{G}_{2\bar{1}}) = 0. \quad (44)$$

(Note that here we used the fact that we had chosen the complex coordinates for the initial hyperkähler base such that the volume factor is unit, *i.e.* $V = 1$.)

It is clear that there always exist such ω^1 and ω^2 that satisfy (43). For example, we can take $z^1 = \int H_1(\omega^i) d\omega^1$ and $z^2 = \omega^2$. Also note that

$$\begin{aligned} A &= -\frac{i}{4}(\tilde{\partial} - \bar{\partial})\partial_y G(y, \omega^i, \bar{\omega}^i) = -\frac{i}{4}(\tilde{\partial} - \bar{\partial})\partial_y (G(y, \omega^i, \bar{\omega}^i) + K_0(\omega^i, \bar{\omega}^i)) \\ &= -\frac{i}{4}(\tilde{\partial} - \bar{\partial})\partial_y (\tilde{G}(y, z^i, \bar{z}^i) + K_0(z^i, \bar{z}^i)) = -\frac{i}{4}(\tilde{\partial} - \bar{\partial})\partial_y \tilde{G}(y, z^i, \bar{z}^i). \end{aligned} \quad (45)$$

Thus the above transformation is a gauge transformation that preserves our initial ansatz. Hence we can set $g = 1$ by this gauge transformation. It follows from (37) that $\kappa_0 = 0$. Now, we have demonstrated that the system with $\kappa_1 = 0$ is determined solely by the following basic equation for G

$$\partial_y^2 G + 4(1 + G_{1\bar{1}} + G_{2\bar{2}} + G_{1\bar{1}}G_{2\bar{2}} - G_{1\bar{2}}G_{2\bar{1}}) = 0. \quad (46)$$

It can be regarded as the $\kappa_1 \rightarrow 0$ limit of (36) if we recover the κ_1 therein. (In the special case when the base space is the flat $\mathbb{R}^4$, the equation (46) was also obtained in [40], but with a numerical error. The factor “8” in equation (2.51) of [40] should be “16” instead.)

To summarize, we find that the Calabi-Yau metrics depend on a discrete choice of the κ function. One is that $\kappa = \alpha$, in which case the solution is completely determined by one basic equation for G , given by (36). The other is that $\kappa = 0$, in which case the solution is completely determined by the basic equation (46).

3 The $\mathbb{R}^4$ base

Having obtained the general formalism for constructing the CY3 metrics from any hyperkähler metric in four dimensions, we consider explicit examples in this and the next sections. Note that all the hyperkähler bases are related by a modification of Kähler potential. Therefore, they are equivalent to each other in our previously general construction. However, since the general basic equation is impossible to solve fully, different choices of hyperkähler base will give different result when we construct the explicit metric in certain simplified ansatz. The simplest hyperkähler space is the Euclidean space $\mathbb{R}^4$. An obvious

choice is to use the complex coordinates (z_1, z_2) directly, and the corresponding complex vielbein is given by

$$\tilde{e}^1 = dz_1, \quad \tilde{e}^2 = dz_2. \quad (47)$$

Alternatively, one can write the $\mathbb{R}^4$ in terms of the spherical-polar coordinates, *i.e.*

$$ds^2 = dr^2 + \frac{1}{4}r^2(\sigma_1^2 + \sigma_2^2 + \sigma_3^2), \quad (48)$$

with the following choice of the complex vielbein

$$\tilde{e}^1 = \tilde{e}^1 + i\tilde{e}^2 = dr + \frac{i}{2}r\sigma_3, \quad \tilde{e}^2 = \tilde{e}^3 + i\tilde{e}^4 = \frac{1}{2}r(\sigma_1 + i\sigma_2). \quad (49)$$

Here, we define the $SU(2)$ Maurer-Cartan forms by

$$\begin{aligned} \sigma_1 &= \sin\psi d\theta - \sin\theta \cos\psi d\phi, \\ \sigma_2 &= -\cos\psi d\theta - \sin\theta \sin\psi d\phi, \\ \sigma_3 &= d\psi + \cos\theta d\phi. \end{aligned} \quad (50)$$

From the relation

$$z_1 = r \cos \frac{\theta}{2} \exp\left(\frac{i(\psi + \phi)}{2}\right), \quad z_2 = r \sin \frac{\theta}{2} \exp\left(\frac{i(\psi - \phi)}{2}\right), \quad (51)$$

one can show that

$$\tilde{e}_a^\mu \partial_\mu z_i = 0. \quad (52)$$

Therefore, the two choices of the complex vielbein (47) and (51) are compatible.

For our purpose, we find that the vielbein (51) is more useful for simplifying equations under the isometry group of the S^3 level surfaces. Under this choice, the general ansatz is

$$\begin{aligned} ds^2 &= (1 + G_{1\bar{1}})(dr^2 + \frac{1}{4}r^2\sigma_3^2) + \frac{1}{4}(1 + G_{2\bar{2}})r^2(\sigma_1^2 + \sigma_2^2) \\ &\quad + (G_{1\bar{2}} + G_{2\bar{1}})(\frac{1}{2}r dr \sigma_1 + \frac{1}{4}r^2 \sigma_3 \sigma_2) - i(G_{1\bar{2}} - G_{2\bar{1}})(\frac{1}{2}r dr \sigma_2 - \frac{1}{4}r^2 \sigma_3 \sigma_1) \\ &\quad + h^2 dy^2 + \frac{1}{h^2}(d\alpha + A)^2. \end{aligned} \quad (53)$$

The inverse complex vielbein is given by

$$\tilde{e}_1 = \frac{1}{2}(\tilde{E}_1 - i\tilde{E}_2), \quad \tilde{e}_2 = \frac{1}{2}(\tilde{E}_3 - i\tilde{E}_4), \quad (54)$$

where

$$\begin{aligned} \tilde{E}_1 &= E_r, \\ \tilde{E}_2 &= \frac{2}{r} E_\psi, \end{aligned}$$

$$\begin{aligned}\tilde{E}_3 &= \frac{2}{r}(\sin \psi E_\theta - \frac{\cos \psi}{\sin \theta} E_\phi + \frac{\cos \theta \cos \psi}{\sin \theta} E_\psi), \\ \tilde{E}_4 &= \frac{2}{r}(-\cos \psi E_\theta - \frac{\sin \psi}{\sin \theta} E_\phi + \frac{\cos \theta \sin \psi}{\sin \theta} E_\psi).\end{aligned}\quad (55)$$

Note that $\Gamma_{i\bar{j}}^{\bar{k}} = \Gamma_{i\bar{j}}^k = 0$ on the Kähler manifold, thus we have

$$\begin{aligned}G_{a\bar{b}} &= \tilde{\epsilon}_a^i \tilde{\epsilon}_b^{\bar{j}} \partial_{z^i} \partial_{\bar{z}^j} G = \tilde{\epsilon}_a^i \tilde{\epsilon}_b^{\bar{j}} \nabla_{z^i} \nabla_{\bar{z}^j} G \\ &= \tilde{\epsilon}_a^\mu \tilde{\epsilon}_b^\nu \nabla_\mu \nabla_\nu G = \tilde{\epsilon}_a^\mu \tilde{\epsilon}_b^\nu (\partial_\mu \partial_\nu - \Gamma_{\mu\nu}^\lambda \partial_\lambda) G.\end{aligned}\quad (56)$$

The explicit form of the $G_{a\bar{b}}$ is presented in (110). The 1-form connection A is given by

$$\begin{aligned}A &= -\frac{i}{4}(\tilde{\partial} - \bar{\partial})\partial_y G = -\frac{i}{4}(\tilde{\epsilon}^a \tilde{\epsilon}_a^\mu - \bar{\epsilon}^a \bar{\epsilon}_a^\mu) \partial_\mu \partial_y G \\ &= \frac{\partial_r \partial_y G}{8} r \sigma_3 - \frac{\partial_\theta \partial_y G}{4} \sin \theta d\phi + \frac{\partial_\phi \partial_y G}{4 \sin \theta} d\theta - \frac{\partial_\psi \partial_y G}{2r} dr - \frac{\partial_\psi \partial_y G}{4 \sin \theta} \cos \theta d\theta.\end{aligned}\quad (57)$$

Since the general equations are rather complicated, we shall further suppose that the functions G and h depend on the coordinates (r, y) only as in [40]. In such a radial ansatz, the resulting metric has the $SU(2) \times U(1)^2$ isometry. Note that we have $G_{1\bar{2}} = 0$ when $G = G(r, y)$. Thus the metric ansatz is reduced to the following form

$$ds^2 = f_1(dr^2 + \frac{1}{4}r^2\sigma_3^2) + \frac{1}{4}f_2r^2(\sigma_1^2 + \sigma_2^2) + \frac{f_1 f_2}{g^2} dy^2 + \frac{g^2}{f_1 f_2} (d\alpha + f_3 \sigma_3)^2. \quad (58)$$

where

$$\begin{aligned}f_2 &\equiv 1 + \frac{1}{2r} \partial_r G, \quad g \equiv f h^{-1} = \sqrt{f_1 f_2} h^{-1}, \\ f_1 &\equiv 1 + \frac{1}{4} \partial_r^2 G + \frac{1}{4r} \partial_r G = \frac{1}{2r} \partial_r (r^2 f_2), \\ f_3 &\equiv \frac{r}{8} \partial_r \partial_y G = \frac{1}{4} \partial_y (r^2 f_2).\end{aligned}\quad (59)$$

3.1 Case I: $\kappa_1 = 1$

In this case, we have

$$g = \exp\left(-\frac{1}{4} \partial_y G\right), \quad (60)$$

and the system is determined solely by the basic equation

$$\partial_y \left[\exp\left(-\frac{1}{2} \partial_y G\right) \right] = \frac{1}{2r^3} \partial_r \left[r^4 \left(1 + \frac{1}{2r} \partial_r G \right)^2 \right]. \quad (61)$$

It is likely difficult to solve this equation fully. We obtain two special solutions: one is just a direct product of Eguchi-Hanson instanton and $\mathbb{R}^2$; the other is given by

$$ds^2 = \frac{dy^2}{W} + \frac{1}{4} W y^2 (d\alpha - r^2 \sigma_3)^2 + y^2 \left(\frac{dr^2}{V} + \frac{1}{4} V r^2 \sigma_3^2 + \frac{1}{4} r^2 (\sigma_1^2 + \sigma_2^2) \right), \quad (62)$$

where

$$W = 1 - \frac{a}{y^6}, \quad V = 1 - r^2 - \frac{b}{r^4}. \quad (63)$$

The detailed derivation can be found in appendix B.1. The metric (62) with $b = 0$ was known in [17], describing a higher dimensional generalization of Eguchi-Hanson instanton, with $\mathbb{R}^2 \times \mathbb{CP}^2$ topology and an asymptotic $\mathbb{R}^6/Z_3$. For $a = 0$, the metric is a cone of $Y^{p,q}$. The general solution describes a resolution of the $Y^{p,q}$ cone, and the detailed global analysis can be found in [8, 9, 10].

It should be emphasized that we have obtained only a special solution to the basic equation (61). It would be interesting to find the general solutions and examine the corresponding metrics.

3.2 Case II: $\kappa_1 = 0$

In this case, we may use f_2 instead of G as a basic function. Then the basic equation (46) becomes

$$\partial_y^2 f_2 + \frac{1}{2r} \partial_r \left(\frac{1}{r^3} \partial_r (r^4 f_2^2) \right) = 0. \quad (64)$$

(This case was discussed in [40]. However, there is an error in the equation (2.52) for G . The constant factor “8” should be “16” instead. This error propagates to the later metric results.)

3.2.1 Some special solutions

One way to solve (64) is to consider the following ansatz

$$f_2 = u(y) r^2 + \xi(y), \quad (65)$$

then we find

$$f_1 = 2 u r^2 + \xi, \quad f_3 = \frac{1}{4} (u_y r^2 + \xi_y) r^2. \quad (66)$$

The functions u and ξ satisfy

$$u_{yy} = -16 u^2, \quad \xi_{yy} = -12 u \xi. \quad (67)$$

An immediate solution is the degenerate case

$$u(y) = 0, \quad \xi(y) = c_1 y + c_2. \quad (68)$$

Beside this case, note that

$$(u_y)^2 = -\frac{32}{3} (u^3 - c^3), \quad (69)$$

$$dy^2 = \frac{3du^2}{32(c^3 - u^3)}, \quad (70)$$

where c is an integration constant. It would be better if we use u instead of y as the coordinate. This implies that ξ satisfies

$$8(c^3 - u^3) \xi_{uu} - 12u^2 \xi_u + 9u \xi = 0. \quad (71)$$

The exact solution for $\xi(u)$ is

$$\begin{aligned} \xi(u) = & {}_2F_1\left(\frac{1}{12}(1 - \sqrt{19}), \frac{1}{12}(1 + \sqrt{19}); \frac{2}{3}; \frac{u^3}{c^3}\right) C_1 \\ & + {}_2F_1\left(\frac{1}{12}(5 - \sqrt{19}), \frac{1}{12}(5 + \sqrt{19}); \frac{4}{3}; \frac{u^3}{c^3}\right) u C_2 \end{aligned} \quad (72)$$

where ${}_2F_1$ is the hypergeometric function. When $c = 0$, the solution takes the simple form

$$\xi(u) = |u|^{-\frac{1}{4} - \frac{\sqrt{19}}{4}} C_1 + |u|^{-\frac{1}{4} + \frac{\sqrt{19}}{4}} C_2. \quad (73)$$

Now the metric becomes

$$\begin{aligned} ds_6^2 = & (2u r^2 + \xi) (dr^2 + \frac{1}{4} r^2 \sigma_3^2) + \frac{1}{4} (u r^2 + \xi) r^2 (\sigma_1^2 + \sigma_2^2) \\ & + \frac{3(2u r^2 + \xi)(u r^2 + \xi)}{32(c^3 - u^3)} du^2 \\ & + \frac{1}{(2u r^2 + \xi)(u r^2 + \xi)} \left(d\alpha + \sqrt{\frac{2}{3}(c^3 - u^3)} (r^2 + \xi_u) r^2 \sigma_3 \right)^2. \end{aligned} \quad (74)$$

We must have $u < c$ to keep the metric real. If $u > 0$, the range of r is $(0, \infty)$. If $u < 0$, then we must have $\xi > 0$, and the range of r is constrained. Especially, when $c \leq 0$, the $r = \infty$ region is not reachable. The metrics have no asymptotic cone over Einstein-Sasaki spaces and they develop a power-law curvature singularity when $f_1 f_2 = 0$. This is rather different from the case of $\kappa_1 = 1$, where the non-vanishing g in the metric (58) allows non-singular collapsing of the cycles.

3.2.2 Separation of variables

We can also solve the equation (64) by separation of variables, namely $f_2 = u(y)\zeta(r)$. Substituting this ansatz to (64), we have

$$u_{yy} + \frac{3}{2}k u^2 = 0, \quad \frac{1}{r} \partial_r (r^{-3} \partial_r (r^4 \zeta^2)) + 3k \zeta = 0, \quad (75)$$

where k is an arbitrary constant. The first equation implies that

$$dy^2 = \frac{du^2}{k(u^3 - c^3)}. \quad (76)$$

The solution for the second equation clearly exists, although there appears to have no explicit analytical form, except for the case with $k = 0$, for which, the solution for f_2 is given by (82).

3.2.3 The formal general solution

Expanding f_2 by Taylor series of y

$$f_2(r, y) = \sum_{n=0}^{\infty} u_n(r) y^n, \quad (77)$$

Substituting this into (64), we find the recursion relation

$$\frac{1}{2r} \partial_r \left(\frac{1}{r^3} \partial_r \left(\sum_{q=0}^n u_q u_{n-q} r^4 \right) \right) + (n+2)(n+1) u_{n+2} = 0. \quad (78)$$

Given the two arbitrarily functions $u_0(r)$ and $u_1(r)$, we can determine all the u_n for $n \geq 2$. The general solution of our system can thus be written formally by the Taylor series of y . If we restrict that both u_0 and u_1 are in the form $u r^2 + \xi_0$, we find that all the u_n 's are in the form $u r^2 + \xi$. It is consistent with the solution we obtained previously.

If we require that there be a maximum $n_{max} = N$ for no vanishing u_n , we shall obtain polynomial solutions on y . There will be $2N + 1$ constraint equations for an N -th order polynomial solution in general except for $N = 0$.

$N = 0$: The only equation in this case is

$$\frac{1}{2r} \partial_r \left(\frac{1}{r^3} \partial_r (u_0^2 r^4) \right) = 0. \quad (79)$$

We find that the solution is

$$f_2 = \sqrt{1 - \frac{a^4}{r^4}} b_0 \quad (80)$$

The corresponding metric is nothing but a direct product of the Eughchi-Hanson instanton and the $\mathbb{R}^2$.

$N = 1$: The equations in this case are

$$\frac{1}{2r} \partial_r \left(\frac{1}{r^3} \partial_r (u_0^2 r^4) \right) = 0, \quad \frac{1}{r} \partial_r \left(\frac{1}{r^3} \partial_r (u_0 u_1 r^4) \right) = 0, \quad \frac{1}{2r} \partial_r \left(\frac{1}{r^3} \partial_r (u_1^2 r^4) \right) = 0. \quad (81)$$

the solution is

$$f_2 = \sqrt{1 - \frac{a^4}{r^4}} (b_0 + b_1 y). \quad (82)$$

$N = 2$: The equations in this case are

$$\begin{aligned} \frac{1}{2r} \partial_r \left(\frac{1}{r^3} \partial_r (u_0^2 r^4) \right) + 2u_2 &= 0, & \frac{1}{r} \partial_r \left(\frac{1}{r^3} \partial_r (u_0 u_1 r^4) \right) &= 0, \\ \frac{1}{2r} \partial_r \left(\frac{1}{r^3} \partial_r ((2u_0 u_2 + u_1^2) r^4) \right) &= 0, & \frac{1}{r} \partial_r \left(\frac{1}{r^3} \partial_r (u_1 u_2 r^4) \right) &= 0, \end{aligned}$$

$$\frac{1}{2r} \partial_r \left(\frac{1}{r^3} \partial_r (u_2^2 r^4) \right) = 0. \quad (83)$$

We find that there is no consistent solution since the solution for the last three equations is in contradiction with the first equation. Similarly, it can be shown that there is no consistent solutions for all finite $N \geq 2$ since the solution for the last $N + 1$ equations are in contradiction with the $(N - 1)$ 'th equation.

4 The triaxial base

In this section, we consider the triaxial hyperkähler metric with the $SU(2)$ isometry. The metric is of the form

$$ds_4^2 = d\rho^2 + a_1^2 \sigma_1^2 + a_2^2 \sigma_2^2 + a_3^2 \sigma_3^2, \quad (84)$$

where a_i 's are functions of ρ only, and they satisfy the first order equations

$$\dot{a}_1 = \frac{a_2^2 + a_3^2 - a_1^2}{2a_2 a_3}, \quad \dot{a}_2 = \frac{a_1^2 + a_3^2 - a_2^2}{2a_1 a_3}, \quad \dot{a}_3 = \frac{a_1^2 + a_2^2 - a_3^2}{2a_1 a_2}. \quad (85)$$

where a dot denotes a derivative with respect to ρ . We find that the most general solution can be written as follows

$$ds_4^2 = \sqrt{\frac{1}{W \tilde{W}}} dr^2 + \frac{1}{4} r^2 (\sqrt{W \tilde{W}} \sigma_1^2 + \sqrt{\frac{\tilde{W}}{W}} \sigma_2^2 + \sqrt{\frac{W}{\tilde{W}}} \sigma_3^2), \quad (86)$$

where

$$W = 1 - \frac{a^4}{r^4}, \quad \tilde{W} = 1 - \frac{b^4}{r^4}. \quad (87)$$

It is of interest to note that one can also introduce a cosmological constant and construct the triaxial Einstein-Kähler spaces. The three first-order equations were obtained in [42]. Only two exact solutions were known: one describes an $\mathbb{CP}^2$ [43] and the other, a direct product $S^2 \times S^2$ [44, 34]. The metrics are

$$\begin{aligned} ds_{\mathbb{CP}^2}^2 &= d\rho^2 + \sin^2 \rho \sigma_1^2 + \cos^2 \rho \sigma_2^2 + \cos^2 2\rho \sigma_3^2, \\ ds_{S^2 \times S^2}^2 &= d\rho^2 + \sin^2 \rho \sigma_1^2 + \sigma_2^2 + \cos^2 \rho \sigma_3^2, \end{aligned} \quad (88)$$

For the generic choice of constant parameters a and b , the metric (86) contains a naked power-law singularity at $r = \max(a, b)$. When $b = a$, the metric reduces to the Eguchi-Hanson instanton, given by

$$ds_4^2 = \frac{dr^2}{W} + \frac{1}{4} W r^2 \sigma_3^2 + \frac{1}{4} r^2 (\sigma_1^2 + \sigma_2^2). \quad (89)$$

We shall first consider this biaxial hyperkähler base. One way to choose the complex vielbein is

$$\tilde{\epsilon}^1 = W^{-\frac{1}{2}} dr + \frac{i}{2} W^{\frac{1}{2}} r \sigma_3, \quad \tilde{\epsilon}^2 = \frac{1}{2} r (\sigma_1 + i \sigma_2). \quad (90)$$

It is easy to see that such a choice will lead to the same radial ansatz as in the $\mathbb{R}^4$ case. Fortunately, there are three possible Kähler structures for a hyperkähler space. For the flat base, all the three choices give same ansatz. For the Eguchi-Hanson base, the remaining two choices are equivalent to each other but inequivalent to (90). The corresponding complex vielbein is given by

$$\tilde{\epsilon}^1 = \tilde{e}^1 + i \tilde{e}^2 = W^{-\frac{1}{2}} dr + \frac{i}{2} r \sigma_3, \quad \tilde{\epsilon}^2 = \tilde{e}^3 + i \tilde{e}^4 = \frac{1}{2} r (W^{\frac{1}{2}} \sigma_1 + i \sigma_2), \quad (91)$$

where we have permuted the σ_i 's for convenience. Correspondingly, the inverse complex vielbein is given by

$$\tilde{\epsilon}_1 = \frac{1}{2}(\tilde{E}_1 - i \tilde{E}_2), \quad \tilde{\epsilon}_2 = \frac{1}{2}(\tilde{E}_3 - i \tilde{E}_4), \quad (92)$$

where

$$\begin{aligned} \tilde{E}_1 &= W^{\frac{1}{2}} E_r, \\ \tilde{E}_2 &= \frac{2}{r} E_\psi, \\ \tilde{E}_3 &= \frac{2}{r W^{\frac{1}{2}}} (\sin \psi E_\theta - \frac{\cos \psi}{\sin \theta} E_\phi + \frac{\cos \theta \cos \psi}{\sin \theta} E_\psi), \\ \tilde{E}_4 &= \frac{2}{r} (-\cos \psi E_\theta - \frac{\sin \psi}{\sin \theta} E_\phi + \frac{\cos \theta \sin \psi}{\sin \theta} E_\psi). \end{aligned} \quad (93)$$

It follows from the relation (56) that $G_{a\bar{b}}$ can be obtained. The result is presented in (111). The 1-form A is given by

$$\begin{aligned} A &= -\frac{i}{4}(\tilde{\partial} - \bar{\partial})\partial_y G = -\frac{i}{4}(\tilde{\epsilon}^a \tilde{\epsilon}_a^\mu - \bar{\tilde{\epsilon}}^a \bar{\tilde{\epsilon}}_a^\mu) \partial_\mu \partial_y G \\ &= \frac{\partial_r \partial_y G}{8} r W^{\frac{1}{2}} \sigma_3 + \frac{\partial_\theta \partial_y G}{4} (W^{\frac{1}{2}} \cos \psi \sigma_1 + W^{-\frac{1}{2}} \sin \psi \sigma_2) \\ &\quad + \frac{\partial_\phi \partial_y G}{4 \sin \theta} (W^{\frac{1}{2}} \sin \psi \sigma_1 - W^{-\frac{1}{2}} \cos \psi \sigma_2) \\ &\quad - \frac{\partial_\psi \partial_y G}{4} \left(\frac{2}{r W^{\frac{1}{2}}} dr + W^{\frac{1}{2}} \cot \theta \sin \psi \sigma_1 - W^{-\frac{1}{2}} \cot \theta \cos \psi \sigma_2 \right). \end{aligned} \quad (94)$$

The metric in the radial ansatz is then given by the following form

$$ds^2 = f_1 \left(\frac{dr^2}{W} + \frac{1}{4} r^2 \sigma_3^2 \right) + \frac{1}{4} f_2 r^2 (W \sigma_1^2 + \sigma_2^2) + \frac{f_1 f_2}{g^2} dy^2 + \frac{g^2}{f_1 f_2} (d\alpha + f_3 \sigma_3)^2. \quad (95)$$

where

$$f_2 = 1 + \frac{1}{2r} \partial_r G, \quad g = f h^{-1} = \sqrt{f_1 f_2} h^{-1},$$

$$\begin{aligned}
f_1 &= 1 + \frac{1}{4}W \partial_r^2 G + \frac{1}{4r} \left(1 + \frac{a^4}{r^4}\right) \partial_r G = \frac{W^{\frac{1}{2}}}{2r} \partial_r \left(r^2 W^{\frac{1}{2}} f_2\right), \\
f_3 &= \frac{r W^{\frac{1}{2}}}{8} \partial_r \partial_y G = \frac{1}{4} r^2 W^{\frac{1}{2}} \partial_y f_2.
\end{aligned} \tag{96}$$

The metric has the isometry of $SU(2) \times U(1)$.

4.1 Case I: $\kappa_1 = 1$

In this case, we have

$$g = \exp \left(-\frac{1}{4} \partial_y G \right), \tag{97}$$

and the system is determined solely by the basic equation

$$\partial_y \left[\exp \left(-\frac{1}{2} \partial_y G \right) \right] = \frac{1}{2r^3} \partial_r \left[(r^4 - a^4) \left(1 + \frac{1}{2r} \partial_r G \right)^2 \right]. \tag{98}$$

We obtain some special solutions. One describes an $\mathbb{R}^2 \times \mathbb{CP}^2$ instanton that is asymptotic to the $\mathbb{R}^6/Z_3$. Another describes an $\mathbb{R}^2 \times S^2 \times S^2$ instanton that is asymptotic to the cone over $T^{1,1}/Z_2$. The details are presented in appendix B.2.

4.2 Case II: $\kappa_1 = 0$

In this case, we may use f_2 instead of G as the basic function. Then the basic equation (46) becomes

$$\partial_y^2 f_2 + \frac{1}{2r} \partial_r \left(\frac{1}{r^3} \partial_r ((r^4 - a^4) f_2^2) \right) = 0. \tag{99}$$

4.2.1 Some special solution

If f_2 depends only on y , the solution is simply $f_2 = c_1 y + c_2$ and the metric is

$$\begin{aligned}
ds^2 &= (c_1 y + c_2) \left(\frac{dr^2}{W} + \frac{1}{4} r^2 \sigma_3^2 \right) + \frac{1}{4} (c_1 y + c_2) r^2 (W \sigma_1^2 + \sigma_2^2) \\
&\quad + (c_1 y + c_2)^2 dy^2 + \frac{1}{(c_1 y + c_2)^2} \left(d\alpha + \frac{c_1}{4} r^2 W^{\frac{1}{2}} \sigma_3 \right)^2.
\end{aligned} \tag{100}$$

If $f_2 = u(y) r^2$, the basic function is simplified to

$$u_{yy} = -16u^2. \tag{101}$$

It implies

$$dy^2 = \frac{3du^2}{32(c^3 - u^3)}. \tag{102}$$

Then we can use u instead of y as the coordinate, and the metric is given by

$$ds_6^2 = (1+W)u r^2 \left(\frac{dr^2}{W} + \frac{1}{4} r^2 \sigma_3^2 \right) + \frac{1}{4} u r^4 (W \sigma_1^2 + \sigma_2^2) + \frac{3(1+W)u^2 r^4}{32(c^3 - u^3)} du^2 + \frac{1}{(1+W)u^2 r^4} \left(d\alpha + \sqrt{\frac{2}{3}(c^3 - u^3)} r^4 W^{\frac{1}{2}} \sigma_3 \right)^2. \quad (103)$$

4.2.2 The formal general solution

The general solution can be expressed formally by Taylor series of y as previously. Expanding f_2 as

$$f_2(r, y) = \sum_{n=0}^{\infty} u_n(r) y^n, \quad (104)$$

we find the recursion relations

$$\frac{1}{2r} \partial_r \left(\frac{1}{r^3} \partial_r \left(\sum_{q=0}^n u_q u_{n-q} (r^4 - a^4) \right) \right) + (n+2)(n+1)u_{n+2} = 0. \quad (105)$$

Given the two arbitrary functions $u_0(r)$ and $u_1(r)$, we can determine all the $u_n (n > 1)$'s by the above recursion relations. Then the general solution of our system can be written formally by the Taylor series of y . If we restrict that both u_0 and u_1 are proportion to r^2 , we find that all the u_n 's are proportion to r^2 by the recursion relations. It is consistent with the previous result. If we require there is a maximum $n_{max} = N$ for no vanishing u_n , we shall obtain polynomial solutions on y . There are $2N + 1$ constraint equations for an N -th order polynomial solution. We now examine these equations as follows.

$N = 0$: The only equation in this case is

$$\frac{1}{2r} \partial_r \left(\frac{1}{r^3} \partial_r (u_0^2 (r^4 - a^4)) \right) = 0. \quad (106)$$

We find that the solution is

$$f_2 = \lambda \sqrt{\frac{r^4 - b^4}{r^4 - a^4}}. \quad (107)$$

It gives rise to a direct product of the triaxial metric (86) and the $\mathbb{R}^2$.

$N = 1$: The equations in this case are

$$\begin{aligned} \frac{1}{2r} \partial_r \left(\frac{1}{r^3} \partial_r (u_0^2 (r^4 - a^4)) \right) &= 0, & \frac{1}{r} \partial_r \left(\frac{1}{r^3} \partial_r (u_0 u_1 (r^4 - a^4)) \right) &= 0, \\ \frac{1}{2r} \partial_r \left(\frac{1}{r^3} \partial_r (u_1^2 (r^4 - a^4)) \right) &= 0. \end{aligned} \quad (108)$$

The solution is

$$f_2 = \sqrt{\frac{r^4 - b^4}{r^4 - a^4}} (\lambda_0 + \lambda_1 y) \quad (109)$$

As in the flat base case, we find again that there are no finite polynomial solutions of the y coordinate, for any order $N \geq 2$.

5 Conclusions

In this paper, we examine the metric construction for Calabi-Yau 3-folds proposed in [40]. The essence of the construction is to add a complex line bundle over a four dimensional hyperkähler structure, with a simple deformation where the four-dimensional Kähler potential is modified. The resulting metric ansatz has at least one Killing direction. It was demonstrated for the $\mathbb{R}^4$ base that the condition for the CY3 metrics could be indeed satisfied.

We extend the construction and obtain the general formalism for a generic hyperkähler base. Furthermore, we find that the ansatz for the holomorphic $(3,0)$ -form should be generalized to have a $U(1)$ factor. This allows us to construct the CY3 metrics that are asymptotic to the cones of Einstein-Sasaki spaces. There can be a discrete choice for the $U(1)$ factor. One is that it depends on the fibre $U(1)$ coordinate only, and consequently the equations of motion are reduced to one differential equation on the modified Kähler potential. The other is that the $U(1)$ factor vanishes. In this case, the metrics are determined by a differential equation that is the singular limit of the previous one.

We then construct explicit metrics with two examples of the hyperkähler bases. One is the $\mathbb{R}^4$, and the other is the triaxial metric with $SU(2)$ isometries. In both cases, we obtain explicit cohomogeneity-2 metrics. With the $U(1)$ factor, we obtain a general class of solutions that describe a resolution of the cone over $Y^{p,q}$ spaces. For the case with vanishing $U(1)$ factor, we obtain singular metrics with no asymptotically conical region. The solutions are governed by two arbitrary functions of the radial variable of the hyperkähler spaces.

The general construction we have obtained allows one to construct a wide class of CY3 metrics with at least one Killing direction. It is of great interest to investigate whether new complete metrics on the non-compact manifolds can arise.

A Explicit $G_{a\bar{b}}$

In this appendix we give the explicit expressions for the $G_{a\bar{b}}$ defined by (56). For the $\mathbb{R}^4$ base discussed in section 3, we find that they are given by

$$\begin{aligned} G_{1\bar{1}} &= \frac{1}{4} \partial_r^2 G + \frac{1}{r^2} \partial_\psi^2 G + \frac{1}{4r} \partial_r G, \\ G_{2\bar{2}} &= \frac{1}{r^2} \partial_\theta^2 G + \frac{1}{r^2 \sin^2 \theta} \partial_\phi^2 G - \frac{2 \cos \theta}{r^2 \sin^2 \theta} \partial_\phi \partial_\psi G + \frac{\cos^2 \theta}{r^2 \sin^2 \theta} \partial_\psi^2 G \\ &\quad + \frac{1}{2r} \partial_r G + \frac{\cos \theta}{r^2 \sin \theta} \partial_\theta G, \\ G_{1\bar{2}} &= e^{i\psi} \left[-\frac{1}{2r \sin \theta} \partial_r \partial_\phi G + \frac{\cos \theta}{2r \sin \theta} \partial_r \partial_\psi G - \frac{1}{r^2} \partial_\theta \partial_\psi G \right] \end{aligned}$$

$$\begin{aligned}
& + i \left(-\frac{1}{2r} \partial_r \partial_\theta G + \frac{1}{r^2 \sin \theta} \partial_\phi \partial_\psi G - \frac{\cos \theta}{r^2 \sin \theta} \partial_\psi^2 G \right) \Big] \\
& = (G_{2\bar{1}})^* .
\end{aligned} \tag{110}$$

For the triaxial base discussed in section 4, they are more complicated, given by

$$\begin{aligned}
G_{1\bar{1}} &= \frac{1}{4} \left(1 - \frac{a^4}{r^4} \right) \partial_r^2 G + \frac{1}{r^2} \partial_\psi^2 G + \frac{1}{4r} \left(1 + \frac{a^4}{r^4} \right) \partial_r G , \\
G_{2\bar{2}} &= \frac{r^4 - a^4 \cos^2 \psi}{r^2(r^4 - a^4)} \partial_\theta^2 G - \frac{a^4 \sin 2\psi}{r^2(r^4 - a^4) \sin \theta} \partial_\theta \partial_\phi G + \frac{a^4 \cos \theta \sin 2\psi}{r^2(r^4 - a^4) \sin \theta} \partial_\theta \partial_\psi G \\
&+ \frac{r^4 - a^4 \sin^2 \psi}{r^2(r^4 - a^4) \sin^2 \theta} \partial_\phi^2 G - \frac{2(r^4 - a^4 \sin^2 \psi) \cos \theta}{r^2(r^4 - a^4) \sin^2 \theta} \partial_\phi \partial_\psi G \\
&+ \frac{(r^4 - a^4 \sin^2 \psi) \cos^2 \theta}{r^2(r^4 - a^4) \sin^2 \theta} \partial_\psi^2 G + \frac{1}{2r} \partial_r G + \frac{(r^4 - a^4 \sin^2 \psi) \cos \theta}{r^2(r^4 - a^4) \sin \theta} \partial_\theta G \\
&+ \frac{a^4 \cos \theta \sin 2\psi}{r^2(r^4 - a^4) \sin^2 \theta} \partial_\phi G - \frac{a^4 (1 + \cos^2 \theta) \sin 2\psi}{2r^2(r^4 - a^4) \sin^2 \theta} \partial_\psi G , \\
G_{1\bar{2}} &= \frac{\sin \psi}{2r} \partial_r \partial_\theta G - \frac{\cos \psi}{2r \sin \theta} \partial_r \partial_\phi G + \frac{\cos \theta \cos \psi}{2r \sin \theta} \partial_r \partial_\psi G - \frac{\cos \psi}{r^2} \partial_\theta \partial_\psi G \\
&- \frac{\sin \psi}{r^2 \sin \theta} \partial_\phi \partial_\psi G + \frac{\cos \theta \sin \psi}{r^2 \sin \theta} \partial_\psi^2 G - \frac{a^4 \sin \psi}{r^2(r^4 - a^4)} \partial_\theta G \\
&+ \frac{a^4 \cos \psi}{r^2(r^4 - a^4) \sin \theta} \partial_\phi G - \frac{a^4 \cos \theta \cos \psi}{r^2(r^4 - a^4) \sin \theta} \partial_\psi G \\
&+ i \left[-\frac{W^{\frac{1}{2}} \cos \psi}{2r} \partial_r \partial_\theta G - \frac{W^{\frac{1}{2}} \sin \psi}{2r \sin \theta} \partial_r \partial_\phi G + \frac{W^{\frac{1}{2}} \cos \theta \sin \psi}{2r \sin \theta} \partial_r \partial_\psi G \right. \\
&- \frac{\sin \psi}{r^2 W^{\frac{1}{2}}} \partial_\theta \partial_\psi G + \frac{\cos \psi}{r^2 W^{\frac{1}{2}} \sin \theta} \partial_\phi \partial_\psi G - \frac{\cos \theta \cos \psi}{r^2 W^{\frac{1}{2}} \sin \theta} \partial_\psi^2 G - \frac{a^4 \cos \psi}{r^6 W^{\frac{1}{2}}} \partial_\theta G \\
&- \frac{a^4 \sin \psi}{r^6 W^{\frac{1}{2}} \sin \theta} \partial_\phi G + \left. \frac{a^4 \cos \theta \sin \psi}{r^6 W^{\frac{1}{2}} \sin \theta} \partial_\psi G \right] = (G_{2\bar{1}})^* .
\end{aligned} \tag{111}$$

B Detailed derivation for the $\kappa = 1$ solutions

B.1 The $\mathbb{R}^4$ base

For the $\kappa = 1$ case, the system is reduced to one basic equation, given by (61). In this appendix, we obtain some special solutions by considering the following ansatz

$$g^2 = u_1(r) (a_3 y^3 + a_2 y^2 + 3a_1^2 y + \tilde{a}_0) . \tag{112}$$

Correspondingly, we have

$$\begin{aligned}
-\frac{1}{2} G &= (2 \log g - 3) y - \sum_{i=1}^3 y_i \log(y - y_i) + u_2(r) \\
&= y \log u_1 + y \log(a_3 y^3 + a_2 y^2 + 3a_1^2 y + \tilde{a}_0) - 3y - \sum_{i=1}^3 y_i \log(y - y_i) + u_2(r) ,
\end{aligned} \tag{113}$$

where y_i 's are the roots of the equation $a_3 y^3 + a_2 y^2 + 3a_1^2 y + \tilde{a}_0 = 0$. Substituting this into the basic equation (61), we have

$$\begin{aligned}\frac{1}{2r^3} \partial_r \left[r^2 \left(\frac{u_1'}{u_1} \right)^2 \right] &= 3a_3 u_1, \\ \frac{1}{2r^3} \partial_r \left[r^2 (u_2' - r) \frac{u_1'}{u_1} \right] &= a_2 u_1, \\ \frac{1}{2r^3} \partial_r \left[r^2 (u_2' - r)^2 \right] &= 3a_1^2 u_1.\end{aligned}\tag{114}$$

If $a_3 = 0$, the consistency requires also $a_2 = 0$. Then we get

$$u_1 = \text{constant}, \quad f_2 = 1 - \frac{u_2'}{r} = a_1 \sqrt{\frac{3u_1}{2} \left(1 + \frac{a}{r^4} \right)}.\tag{115}$$

After absorbing the redundant parameter by rescaling, the metric is given by

$$ds^2 = \frac{1}{\sqrt{1 + \frac{a^4}{r^4}}} (dr^2 + \frac{1}{4} r^2 \sigma_3^2) + \frac{r^2}{4} \sqrt{1 + \frac{a^4}{r^4}} (\sigma_1^2 + \sigma_2^2) + dy^2 + d\alpha^2.\tag{116}$$

Further making the coordinate transformation

$$(r^4 + a^4)^{\frac{1}{4}} \rightarrow r,\tag{117}$$

we get

$$ds^2 = \frac{dr^2}{1 - \frac{a^4}{r^4}} + \frac{r^2}{4} \left(1 - \frac{a^4}{r^4} \right) \sigma_3^2 + \frac{1}{4} r^2 (\sigma_1^2 + \sigma_2^2) + dy^2 + d\alpha^2.\tag{118}$$

This is nothing but a direct product of the Eughchi-Hanson instanton and the $\mathbb{R}^2$.

If $a_3 \neq 0$, we can always set $a_3 = 1$ by the redefinition of u . Then the consistency of the equations implies

$$u_2 = \frac{1}{2} r^2 + a_1 \log u_1, \quad a_2 = 3a_1.\tag{119}$$

Thus

$$\begin{aligned}g^2 &= u_1(r) (y^3 + 3a_1 y^2 + 3a_1^2 y + \tilde{a}_0) = u_1(r) [(y + a_1)^3 + a_0], \\ r^2 f_2 &= r^2 \left(1 + \frac{1}{2r} \partial_r G \right) = -(y + a_1) r \partial_r \log u_1, \\ r^2 f_1 &= \frac{r}{2} \partial_r (r^2 f_2) = -(y + a_1) \frac{r}{2} \partial_r (r \partial_r \log u_1), \\ f_3 &= \frac{1}{4} \partial_y (r^2 f_2) = -\frac{r}{4} \partial_r \log u_1 \\ h^2 &= \frac{f_1 f_2}{g^2} = \frac{(y + a_1)^2 \partial_r (r \partial_r \log u_1) \partial_r \log u_1}{2 [(y + a_1)^3 + a_0] r^2 u_1} = \frac{3(y + a_1)^2}{2 [(y + a_1)^3 + a_0]}.\end{aligned}\tag{120}$$

Obviously, we can set $a_1 = 0$ by a coordinate transformation. Then the metric is given by

$$ds^2 = -\frac{y}{2} \partial_r \rho \left(\frac{dr^2}{r} + \frac{1}{4} r \sigma_3^2 \right) - \frac{y}{4} \rho (\sigma_1^2 + \sigma_2^2)$$

$$+\frac{3y^2}{2(y^3+a_0)}dy^2+\frac{2(y^3+a_0)}{3y^2}(d\alpha-\frac{1}{4}\rho\sigma_3)^2, \quad (121)$$

where

$$\rho = r \partial_r \log u_1. \quad (122)$$

Taking ρ instead of r as the radial coordinate and supposing

$$r \partial_r = V(\rho) \partial_\rho, \quad (123)$$

the metric becomes

$$ds^2 = -\frac{y}{2} \left(\frac{d\rho^2}{V} + \frac{V}{4} \sigma_3^2 \right) - \frac{y}{4} \rho (\sigma_1^2 + \sigma_2^2) \\ + \frac{3y^2}{2(y^3+a_0)} dy^2 + \frac{2(y^3+a_0)}{3y^2} (d\alpha - \frac{1}{4} \rho \sigma_3)^2. \quad (124)$$

The first equation in (114) becomes

$$3u_1 = \frac{V}{2r^4} \partial_\rho(\rho^2) = \frac{\rho V}{r^4}. \quad (125)$$

Thus

$$\rho = V \partial_\rho \log u_1 = \frac{1}{\rho} \partial_\rho(\rho V) - 4. \quad (126)$$

The solution is

$$V = \left(\frac{1}{3} \rho^3 + 2\rho^2 + b_0 \right) \rho^{-1}. \quad (127)$$

Let $\rho \rightarrow -\rho$ for convenience, then the CY3 metric is given by

$$ds^2 = \frac{y}{2} \left(\frac{d\rho^2}{2\rho - \frac{1}{3}\rho^2 + \frac{b_0}{\rho}} + \frac{2\rho - \frac{1}{3}\rho^2 + \frac{b_0}{\rho}}{4} \sigma_3^2 \right) + \frac{y}{4} \rho (\sigma_1^2 + \sigma_2^2) \\ + \frac{3y^2}{2(y^3+a_0)} dy^2 + \frac{2(y^3+a_0)}{3y^2} (d\alpha + \frac{1}{4} \rho \sigma_3)^2. \quad (128)$$

After making some appropriate coordinate transformations, rescaling of the metric and renaming the constants, the metric can be cast into (62).

B.2 The triaxial base

The basic equation for the $\kappa = 1$ solutions is given by (98). Since the structure is quite similar with that of the flat $\mathbb{R}^4$ base, we take the same ansatz as in that case, namely

$$g^2 = u_1(r) (a_3 y^3 + a_2 y^2 + 3a_1^2 y + \tilde{a}_0). \quad (129)$$

Thus we have

$$\begin{aligned}
-\frac{1}{2}G &= (2\log g - 3)y - \sum_{i=1}^3 y_i \log(y - y_i) + u_2(r) \\
&= y \log u_1 + y \log(a_3 y^3 + a_2 y^2 + 3a_1^2 y + \tilde{a}_0) - 3y - \sum_{i=1}^3 y_i \log(y - y_i) + u_2(r), \quad (130)
\end{aligned}$$

where y_i 's are the roots of the equation $a_3 y^3 + a_2 y^2 + 3a_1^2 y + \tilde{a}_0 = 0$. Then, the basic equation implies

$$\begin{aligned}
\frac{1}{2r^3} \partial_r \left[r^2 W \left(\frac{u'_1}{u_1} \right)^2 \right] &= 3a_3 u_1, \\
\frac{1}{2r^3} \partial_r \left[r^2 W (u'_2 - r) \frac{u'_1}{u_1} \right] &= a_2 u_1, \\
\frac{1}{2r^3} \partial_r \left[r^2 W (u'_2 - r)^2 \right] &= 3a_1^2 u_1. \quad (131)
\end{aligned}$$

If $a_3 = 0$, the consistency requires also $a_2 = 0$. The nontrivial solution is given by

$$u_1 = \text{constant}, \quad f_2 = 1 - \frac{u'_2}{r} = a_1 \sqrt{\frac{3u_1}{2}} \sqrt{\frac{r^4 - b^4}{r^4 - a^4}}. \quad (132)$$

After absorbing the redundant parameter by rescaling, the corresponding metric is

$$ds^2 = \sqrt{\frac{1}{W \tilde{W}}} dr^2 + \frac{1}{4} r^2 (\sqrt{W \tilde{W}} \sigma_1^2 + \sqrt{\frac{\tilde{W}}{W}} \sigma_2^2 + \sqrt{\frac{W}{\tilde{W}}} \sigma_3^2) + dy^2 + d\alpha^2. \quad (133)$$

This is just a direct product of the triaxial metric (86) and the $\mathbb{R}^2$. It implies that the radial ansatz for the CY3 metric based on the triaxial hyperkähler base will be equivalent to (95). Therefore, up to coordinate transformations which permute the three σ_i 's, there will be no further radial ansatz coming from the triaxial base.

For non-vanishing a_3 , we can set it to unity by redefinition of u . Then the consistency of the equations implies

$$u_2 = \frac{1}{2} r^2 + a_1 \log u_1, \quad a_2 = 3a_1. \quad (134)$$

Thus

$$\begin{aligned}
g^2 &= u_1(r) (y^3 + 3a_1 y^2 + 3a_1^2 y + \tilde{a}_0) = u_1(r) [(y + a_1)^3 + a_0], \\
r^2 f_2 &= r^2 (1 + \frac{1}{2r} \partial_r G) = -(y + a_1) r \partial_r \log u_1, \\
r^2 f_1 &= \frac{1}{2} r W^{\frac{1}{2}} \partial_r (r^2 W^{\frac{1}{2}} f_2) = -\frac{1}{2} (y + a_1) r W^{\frac{1}{2}} \partial_r (r W^{\frac{1}{2}} \partial_r \log u_1), \\
f_3 &= \frac{1}{4} r^2 W^{\frac{1}{2}} \partial_y f_2 = -\frac{1}{4} r W^{\frac{1}{2}} \partial_r \log u_1 \\
h^2 &= \frac{f_1 f_2}{g^2} = \frac{(y + a_1)^2 W^{\frac{1}{2}} \partial_r (r W^{\frac{1}{2}} \partial_r \log u_1) \partial_r \log u_1}{2 [(y + a_1)^3 + a_0] r^2 u_1} = \frac{3(y + a_1)^2}{2 [(y + a_1)^3 + a_0]}. \quad (135)
\end{aligned}$$

Obviously, we can set $a_1 = 0$ by coordinate transformation. Then the metric is given by

$$ds^2 = -\frac{y}{2} \partial_r \rho \left(\frac{dr^2}{r W^{\frac{1}{2}}} + \frac{1}{4} r W^{\frac{1}{2}} \sigma_3^2 \right) - \frac{y}{4} \rho (W^{\frac{1}{2}} \sigma_1^2 + W^{-\frac{1}{2}} \sigma_2^2) + \frac{3y^2}{2(y^3 + a_0)} dy^2 + \frac{2(y^3 + a_0)}{3y^2} (d\alpha - \frac{1}{4} \rho \sigma_3)^2, \quad (136)$$

where

$$\rho = r W^{\frac{1}{2}} \partial_r \log u_1. \quad (137)$$

Supposing

$$r W^{\frac{1}{2}} \partial_r = \xi \partial_\xi = V(\rho) \partial_\rho, \quad (138)$$

we find

$$\partial_\rho \log \xi = \frac{1}{V}, \quad r^2 = \frac{4\xi^4 + a^4}{4\xi^2}, \quad W^{\frac{1}{2}} = \frac{4\xi^4 - a^4}{4\xi^4 + a^4}. \quad (139)$$

Taking ρ instead of r as the radial coordinate, the metric becomes

$$ds^2 = -\frac{y}{2} \left(\frac{d\rho^2}{V} + \frac{V}{4} \sigma_3^2 \right) - \frac{y}{4} \rho (W^{\frac{1}{2}} \sigma_1^2 + W^{-\frac{1}{2}} \sigma_2^2) + \frac{3y^2}{2(y^3 + a_0)} dy^2 + \frac{2(y^3 + a_0)}{3y^2} (d\alpha - \frac{1}{4} \rho \sigma_3)^2. \quad (140)$$

The first equation in (131) becomes

$$3u_1 = \frac{V}{2r^4 W^{\frac{1}{2}}} \partial_\rho(\rho^2) = \frac{16\xi^4 \rho V}{16\xi^8 - a^8}. \quad (141)$$

Thus

$$\rho = V \partial_\rho \log u_1 = \frac{1}{\rho} \partial_\rho(\rho V) - 4 \frac{16\xi^8 + a^8}{16\xi^8 - a^8}. \quad (142)$$

Then we have

$$\begin{aligned} \frac{16\xi^8}{a^8} &= \frac{\frac{1}{\rho} \partial_\rho(\rho V) - \rho + 4}{\frac{1}{\rho} \partial_\rho(\rho V) - \rho - 4} \\ \Rightarrow \frac{8}{V} &= \partial_\rho \log \left(\frac{\frac{1}{\rho} \partial_\rho(\rho V) - \rho + 4}{\frac{1}{\rho} \partial_\rho(\rho V) - \rho - 4} \right) = - \frac{8 \partial_\rho \left(\frac{1}{\rho} \partial_\rho(\rho V) \right) - 8}{\left(\frac{1}{\rho} \partial_\rho(\rho V) - \rho \right)^2 - 16}. \end{aligned} \quad (143)$$

Let

$$\tilde{V} = \rho V - \frac{1}{3} \rho^3, \quad \tilde{\rho} = \rho^2, \quad (144)$$

the above equation can be rewritten as

$$\left(\tilde{V} + \frac{1}{3} \tilde{\rho}^{\frac{3}{2}} \right) \partial_{\tilde{\rho}}^2 \tilde{V} + (\partial_{\tilde{\rho}} \tilde{V})^2 - 4 = 0. \quad (145)$$

Supposing

$$\tilde{V} = b_3 \tilde{\rho}^{\frac{3}{2}} + b_2 \tilde{\rho} + b_1 \tilde{\rho}^{\frac{1}{2}} + b_0, \quad (146)$$

the corresponding solutions are given by

$$1) \quad \tilde{V} = \pm 2 \tilde{\rho} + b_0, \quad 2) \quad \tilde{V} = -48 \tilde{\rho}^{\frac{1}{2}}, \quad 3) \quad \tilde{V} = -\frac{1}{12} \tilde{\rho}^{\frac{3}{2}} - 16 \tilde{\rho}^{\frac{1}{2}}. \quad (147)$$

The first solution corresponds to the flat base case discussed in the previous section. The second solution gives rise to the metric

$$\begin{aligned} ds^2 = & \frac{y}{2} \left(\frac{3 d\rho^2}{12^2 - \rho^2} + \frac{12^2 - \rho^2}{12} \sigma_3^2 \right) \\ & + \frac{y}{4} \left[\left(12 - \sqrt{12^2 - \rho^2} \right) \sigma_1^2 + \left(12 + \sqrt{12^2 - \rho^2} \right) \sigma_2^2 \right] \\ & + \frac{3 y^2}{2 (y^3 + a_0)} dy^2 + \frac{2 (y^3 + a_0)}{3 y^2} \left(d\alpha - \frac{1}{4} \rho \sigma_3 \right)^2. \end{aligned} \quad (148)$$

With certain coordinate transformation and rescaling of the metric, it can be expressed as

$$ds^2 = \frac{dy^2}{W} + \frac{1}{4} W y^2 (d\alpha - 2 \sin(2\rho) \sigma_3)^2 + y^2 ds_{\mathbb{CP}^2}^2. \quad (149)$$

where W is given by (63) and $ds_{\mathbb{CP}^2}^2$ is the triaxial $\mathbb{CP}^2$ metric given by (88). Thus, the metric describes an $\mathbb{R}^2 \times \mathbb{CP}^2$ instanton that is asymptotic to the $\mathbb{R}^6/Z_3$.

The third solution gives the metric

$$\begin{aligned} ds^2 = & \frac{y}{2} \left(\frac{4 d\rho^2}{8^2 - \rho^2} + \frac{8^2 - \rho^2}{16} \sigma_3^2 \right) + \frac{y}{4} \left(\frac{\rho^2}{8} \sigma_1^2 + 8 \sigma_2^2 \right) \\ & + \frac{3 y^2}{2 (y^3 + a_0)} dy^2 + \frac{2 (y^3 + a_0)}{3 y^2} \left(d\alpha - \frac{1}{4} \rho \sigma_3 \right)^2. \end{aligned} \quad (150)$$

With certain coordinate transformation and rescaling of the metric, it can be expressed as

$$ds^2 = \frac{dy^2}{W} + \frac{1}{4} W y^2 (d\alpha - \frac{4}{3} \sin \rho \sigma_3)^2 + \frac{1}{3} y^2 ds_{S^2 \times S^2}^2. \quad (151)$$

where W is given by (63) and $ds_{S^2 \times S^2}^2$ is the triaxial $S^2 \times S^2$ metric given by (88). Thus, the metric describes an $\mathbb{R}^2 \times S^2 \times S^2$ instanton that is asymptotic to the cone over $T^{1,1}/Z_2$.

Acknowledgement

We are grateful to Chris Pope for useful discussion. Z.L.W. acknowledges the support by grants from the Chinese Academy of Sciences, a grant from 973 Program with grant No: 2007CB815401 and grants from the NSF of China with Grant No:10588503 and 10535060.

References

- [1] E. Calabi, *The space of Kähler metrics*, Proc. Internat. Congress Math. Amsterdam, (1954) 206.
- [2] E. Calabi, *On Kähler manifolds with vanishing canonical class*, Algebraic geometry and topology: A symposium in honor of S. Lefschetz, Princeton University Press, (1957) 78.
- [3] S.T. Yau, *On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation. I*, Communications on Pure and Applied Mathematics 31 (3) (1978) 339.
- [4] P. Candelas and X.C. de la Ossa, *Comments on Conifolds*, Nucl. Phys. B **342**, 246 (1990).
- [5] J.P. Gauntlett, D. Martelli, J. Sparks and D. Waldram, *Sasaki-Einstein metrics on $S^2 \times S^3$* , Adv. Theor. Math. Phys. **8**, 711 (2004) [arXiv:hep-th/0403002].
- [6] M. Cvetič, H. Lü, D.N. Page and C.N. Pope, *New Einstein-Sasaki spaces in five and higher dimensions*, Phys. Rev. Lett. **95**, 071101 (2005) [arXiv:hep-th/0504225].
- [7] M. Cvetič, H. Lü, D.N. Page and C.N. Pope, *New Einstein-Sasaki and Einstein spaces from Kerr-de Sitter*, JHEP **0907**, 082 (2009) [arXiv:hep-th/0505223].
- [8] T. Oota and Y. Yasui, *Explicit toric metric on resolved Calabi-Yau cone*, Phys. Lett. B **639**, 54 (2006) [arXiv:hep-th/0605129].
- [9] H. Lü and C.N. Pope, *Resolutions of cones over Einstein-Sasaki spaces*, Nucl. Phys. B **782**, 171 (2007) [arXiv:hep-th/0605222].
- [10] W. Chen, M. Cvetič, H. Lü, C.N. Pope and J.F. Vazquez-Poritz, *Resolved Calabi-Yau Cones and Flows from L^{abc} Superconformal Field Theories*, Nucl. Phys. B **785**, 74 (2007) [arXiv:hep-th/0701082].
- [11] Y. Hashimoto, M. Sakaguchi and Y. Yasui, *Sasaki-Einstein twist of Kerr-AdS black holes*, Phys. Lett. B **600**, 270 (2004) [arXiv:hep-th/0407114].
- [12] W. Chen, H. Lü and C.N. Pope, *General Kerr-NUT-AdS metrics in all dimensions*, Class. Quant. Grav. **23**, 5323 (2006) [arXiv:hep-th/0604125].

- [13] S.W. Hawking, C.J. Hunter and M. Taylor, *Rotation and the AdS/CFT correspondence*, Phys. Rev. D **59**, 064005 (1999) [arXiv:hep-th/9811056].
- [14] G.W. Gibbons, H. Lü, D.N. Page and C.N. Pope, *The general Kerr-de Sitter metrics in all dimensions*, J. Geom. Phys. **53**, 49 (2005) [arXiv:hep-th/0404008].
- [15] G.W. Gibbons, H. Lü, D.N. Page and C.N. Pope, *Rotating black holes in higher dimensions with a cosmological constant*, Phys. Rev. Lett. **93**, 171102 (2004) [arXiv:hep-th/0409155].
- [16] L. Berard-Bergery, *Quelques exemples de varietes riemanniennes completes non compactes a courbure de Ricci-Positive*, C.R. Acad. Sci., Paris, Ser. I320, 159 (1986).
- [17] D.N. Page and C.N. Pope, *Inhomogeneous Einstein Metrics On Complex Line Bundles*, Class. Quant. Grav. **4**, 213 (1987).
- [18] R.L. Bryant and S. Salamon, *On the construction of some complete metrics with exceptional holonomy*, Duke Math. J. **58**, 829 (1989).
- [19] G.W. Gibbons, D.N. Page and C.N. Pope, *Einstein Metrics on S^3 , $\mathbb{R}^3$ and $\mathbb{R}^4$ Bundles*, Commun. Math. Phys. **127**, 529 (1990).
- [20] M. Cvetič, G.W. Gibbons, H. Lü and C.N. Pope, *New complete non-compact Spin(7) manifolds*, Nucl. Phys. B **620**, 29 (2002) [arXiv:hep-th/0103155].
- [21] A. Brandhuber, J. Gomis, S.S. Gubser and S. Gukov, *Gauge theory at large N and new G_2 holonomy metrics*, Nucl. Phys. B **611**, 179 (2001) [arXiv:hep-th/0106034].
- [22] H. Kanno and Y. Yasui, *On Spin(7) holonomy metric based on $SU(3)/U(1)$* , J. Geom. Phys. **43**, 293 (2002) [arXiv:hep-th/0108226].
- [23] M. Cvetič, G.W. Gibbons, H. Lü and C.N. Pope, *Cohomogeneity one manifolds of Spin(7) and G_2 holonomy*, Phys. Rev. D **65**, 106004 (2002) [arXiv:hep-th/0108245].
- [24] S. Gukov and J. Sparks, *M-theory on Spin(7) manifolds. I*, Nucl. Phys. B **625**, 3 (2002) [arXiv:hep-th/0109025].
- [25] M. Cvetič, G.W. Gibbons, H. Lü and C.N. Pope, *Orientifolds and slumps in G_2 and Spin(7) metrics*, Annals Phys. **310**, 265 (2004) [arXiv:hep-th/0111096].

- [26] G. Curio, B. Kors and D. Lust, *Fluxes and branes in type II vacua and M-theory geometry with G_2 and $Spin(7)$ holonomy*, Nucl. Phys. B **636**, 197 (2002) [arXiv:hep-th/0111165].
- [27] H. Kanno and Y. Yasui, *On $Spin(7)$ holonomy metric based on $SU(3)/U(1)$. II*, J. Geom. Phys. **43**, 310 (2002) [arXiv:hep-th/0111198].
- [28] A. Brandhuber, *G_2 holonomy spaces from invariant three-forms*, Nucl. Phys. B **629**, 393 (2002) [arXiv:hep-th/0112113].
- [29] M. Cvetič, G.W. Gibbons, H. Lu and C.N. Pope, *A G_2 unification of the deformed and resolved conifolds*, Phys. Lett. B **534**, 172 (2002) [arXiv:hep-th/0112138].
- [30] M. Cvetič, G.W. Gibbons, H. Lü and C.N. Pope, *Almost special holonomy in type IIA and M theory*, Nucl. Phys. B **638**, 186 (2002) [arXiv:hep-th/0203060].
- [31] S. Gukov, S.T. Yau and E. Zaslow, *Duality and fibrations on G_2 manifolds*, arXiv:hep-th/0203217.
- [32] Z.W. Chong, M. Cvetič, G.W. Gibbons, H. Lü, C.N. Pope and P. Wagner, *General metrics of $G(2)$ holonomy and contraction limits*, Nucl. Phys. B **638**, 459 (2002) [arXiv:hep-th/0204064].
- [33] H. Lü, *New G_2 metric, D6-branes and lattice universe*, Nucl. Phys. B **642**, 173 (2002) [arXiv:hep-th/0204231].
- [34] M. Cvetič, G.W. Gibbons, H. Lü and C.N. Pope, *Bianchi IX self-dual Einstein metrics and singular G_2 manifolds*, Class. Quant. Grav. **20**, 4239 (2003) [arXiv:hep-th/0206151].
- [35] Z.W. Chong, *General metrics of G_2 and $Spin(7)$ holonomy*, arXiv:hep-th/0411125.
- [36] O.P. Santillan, *New G_2 holonomy metrics, D6 branes with inherent $U(1) \times U(1)$ isometry and gamma-deformations*, Phys. Rev. D **73**, 126011 (2006) [arXiv:hep-th/0605161].
- [37] G. Giribet and O.P. Santillan, *Toric G_2 and $Spin(7)$ holonomy spaces from gravitational instantons and other examples*, Commun. Math. Phys. **275**, 373 (2007) [arXiv:hep-th/0608192].
- [38] O.P. Santillan, *A Kaehler-Einstein inspired ansatz for $spin(7)$ holonomy metrics and its solution*, arXiv:hep-th/0609088.

- [39] S. Salur and O. Santillan, *New Spin(7) holonomy metrics admitting G2 holonomy reductions and M-theory/IIA dualities*, arXiv:0811.4422 [hep-th].
- [40] O.P. Santillan, *New regular non compact Calabi-Yau metrics in $D = 6$* , arXiv:0909.1718 [hep-th].
- [41] A. Fayyazuddin, *Calabi-Yau 3-folds from 2-folds*, Class. Quant. Grav. **24**, 3151 (2007) [arXiv:hep-th/0702135].
- [42] A.S. Dancer and I.A.B. Strachan, *Kahler-Einstein Metrics With $SU(2)$ Action*, Math. Proc. Camb. Phil. Soc. **115**, 513 (1994).
- [43] C. Bouchiat and G.W. Gibbons, *Nonintegrable Quantum Phase In The Evolution Of A Spin 1 System: A Physical Consequence Of The Nontrivial Topology Of The Quantum State Space*, J. Phys. (France) **49**, 187 (1988).
- [44] C.N. Pope, Ph.D. thesis, Cambridge University, 1979.